

GUIDANCE ON THE TIMELINESS OF POST-DISCHARGE CARE FOR ADULTS FOLLOWING ACUTE KIDNEY INJURY





CLINICAL CONTEXT AT POINT OF HOSPITAL DISCHARGE

RCGP AKI TOOLKIT

RCGP INFOGRAPHIC

Post discharge care for adults

following AKI: Top ten tips

Evidence, references

and resources

AKI STAGE

3

AKI

STAGE

2

AKI

STAGE

HEART FAILURE POOR KIDNEY RECOVERY

> CONSIDER CLINICAL **REVIEW BY 3 DAYS**

HEART FAILURE

MODERATE OR GOOD KIDNEY RECOVERY

CONSIDER CLINICAL **REVIEW BY 1-2 WEEKS**

NO OTHER SIGNIFICANT FACTORS (NO HEART FAILURE)

> **POOR** KIDNEY RECOVERY

SIGNIFICANT RISK FACTOR O (NO HEART FAILURE)

MODERATE KIDNEY RECOVERY NO SIGNIFICANT RISK FACTOR

MODERATE KIDNEY RECOVERY

CONSIDER CLINICAL REVIEW BY 1 MONTH

SIGNIFICANT RISK FACTOR O

GOOD KIDNEY RECOVERY

CONSIDER CLINICAL **REVIEW BY 1 MONTH**

CONSIDER U&Es

O SIGNIFICANT RISK FACTORS (IN ADDITION TO HEART FAILURE) PROMPTING EARLIER **REVIEW**

Chronic kidney disease (CKD)

Other cardiovascular risk factors (diabetes, hypertension and established cardiovascular disease)

Markers of vulnerability: recurrent AKI, cancer treatment, sepsis, critical care

Markers of frailty: those defined within the NHS **England toolkit for general** practice in supporting older people living with frailty

OKIDNEY MONITORING FOLLOWING AKI

Why is a test needed? Kidney function has not

Medicines (ACEI/ARB/

restarted/up titrated

MRA/Diuretics) have been

stabilised









BLOOD TEST MONITORING O

CONSIDER U&Es BY 1-2 WEEKS

CONSIDER CLINICAL

REVIEW BY 1-2 WEEKS

BY 1 MONTH

CONSIDER U&Es BY 3 MONTHS O

NO SIGNIFICANT

RISK FACTOR

GOOD KIDNEY RECOVERY

CONSIDER CLINICAL

REVIEW BY 3 MONTHS

URINE ACR \circ

CONSIDER URINE ACR BY 3 MONTHS (

AKI SEVERITY

AKI STAGE 1 SCr ≥1.5 x baseline level (or SCr rise >26 μ mol/L ≤48 hrs) AKI STAGE 2 $SCr \ge 2 \times baseline level$ **AKI STAGE 3** SCr \geq 3 x baseline level (or SCr \geq 1.5 x baseline to >354 μ mol/L)

Based on SCr change known or presumed to have occurred within previous 7 days.

KIDNEY RECOVERY

CONSIDER CLINICAL

REVIEW BY 1-2 WEEKS

Consider the most recent stable creatinine value prior to AKI to determine the degree of kidney recovery. Refer also to the NHS England algorithm for detecting AKI.

GOOD RECOVERY SCr < 25% above baseline

MODERATE RECOVERY SCr >25% & <50% above baseline

POOR RECOVERY SCr > 50%above baseline

ABBREVIATIONS

Albumin/creatinine ratio

AKI

Acute Kidney Injury

Serum creatinine

U&Es

Urea and electrolytes

O CHECK FOR DEVELOPMENT OR PROGRESSION OF CKD

Align with existing reviews to reduce workload and patient burden



This guidance has been developed using established RAND/UCLA methodology.

The guidance is based on consensus on the most appropriate response to a range of scenarios but must not replace clinical judgement based on individual circumstances.

It does not apply to children, young adults (<18y), people with kidney transplants or on dialysis, or people receiving end of life care.



AKI IS ASSOCIATED WITH

- Re-hospitalisation <30 days
- Further AKI
- Development and progression of CKD
- Cardiovascular mortality